

MATHEMATICAL TRIPOS      Part III

---

Wednesday 4 June 2003    1.30 to 4.30

---

PAPER 45

SYMMETRY AND PARTICLE PHYSICS

*Attempt **THREE** questions.*

*There are **four** questions in total.*

*The questions carry equal weight.*

**You may not start to read the questions  
printed on the subsequent pages until  
instructed to do so by the Invigilator.**

**1** What are the electric charges and isospin quantum numbers of the  $u$  and  $d$  quarks?

What are the spins, charges and isospin quantum numbers of the possible particle states which may be formed from three  $u, d$  quarks? What observed baryons, with appropriate spins and isospins, may these particle states be identified with? Describe also the quark, anti-quark meson states formed from the  $u, d$  quarks and their corresponding anti-quarks.

If there is an additional quark with isospin 0 and charge  $\frac{2}{3}$  or  $-\frac{1}{3}$  what additional baryons and mesons may be formed containing one of the new quarks? Determine their isospins, spins and charges.

[No proofs of rules for combining  $SU(2)$  representations need be given but they should be clearly stated. Disregard any orbital angular momentum.]

**2** Explain why irreducible  $SU(3)$  tensors can be restricted to the form  $T_{\beta_1 \dots \beta_l}^{\alpha_1 \dots \alpha_k}$ , for  $\alpha_i, \beta_j \in \{1, 2, 3\}$ , where the indices  $\alpha_1 \dots \alpha_k$  and  $\beta_1 \dots \beta_l$  are completely symmetrised and  $T_{\beta_1 \dots \beta_l}^{\alpha_1 \dots \alpha_k}$  is zero on contraction of any  $\alpha$  index with a  $\beta$  index. Show that the number of independent components is

$$\frac{1}{4}(k+1)(k+2)(l+1)(l+2) - \frac{1}{4}k(k+1)l(l+1) = \frac{1}{2}(k+1)(l+1)(k+l+2).$$

Discuss the tensor product of two irreducible tensors  $T_{\beta}^{\alpha}$  and  $S_{\delta}^{\gamma}$  showing that the dimensions add up as required.

If  $|B_r\rangle$  are baryon octet states,  $r = 1, \dots, 8$  and  $V_i$  is an operator which also transforms as an octet, so that  $[F_i, V_j] = if_{ijk}V_k$  where  $F_i$  are the  $SU(3)$  charges, explain in outline why previous results would suggest that  $\langle B_r | V_i | B_s \rangle$  should depend on two independent parameters for any  $r, s, i$ .

**3** The Lie algebra of  $SU(2)$  can be written in the form

$$[H, E^\pm] = \pm 2E^\pm, \quad [E^+, E^-] = H.$$

Show how a basis for a representation space may be formed by states  $\{|n\rangle\}$ , which are eigenvectors of  $H$ ,  $H|n\rangle = n|n\rangle$ , such that

$$E^-|n\rangle \propto |n-2\rangle \quad \text{or} \quad E^-|n\rangle = 0,$$

starting from a state  $|\bar{n}\rangle$  satisfying  $E^+|\bar{n}\rangle = 0$ . For  $\bar{n} \geq 0$  and  $\bar{n}$  an integer show that a finite dimensional space is obtained.

[You may assume

$$E^+E^-|n\rangle = \frac{1}{4}(\bar{n} + n)(\bar{n} - n + 2)|n\rangle,$$

but indicate how it may be proved by induction.]

What are the possible eigenvalues for  $H$  in the representation and what is its dimension?

A rank 2 Lie algebra has two commuting elements. How are the roots  $\underline{\alpha}$  defined? The simple roots are  $\underline{\alpha}_1, \underline{\alpha}_2$ . Define the Cartan matrix  $[K_{ij}]$  in this case. Show in outline how the Lie algebra can be written in part in the form

$$[H_i, H_j] = 0, \quad [E^+_i, E^-_j] = \delta_{ij}H_j, \quad [H_i, E^\pm_j] = \pm K_{ji}E^\pm_j, \quad \text{no sum on } j.$$

Explain briefly how a representation space with a basis  $\{|n_1, n_2\rangle\}$  where  $H_i|n_1, n_2\rangle = n_i|n_1, n_2\rangle$  may be obtained starting from a state  $|\bar{n}_1, \bar{n}_2\rangle$  satisfying  $E^+_i|\bar{n}_1, \bar{n}_2\rangle = 0$  with  $\bar{n}_i$  integers and  $\bar{n}_i \geq 0$ .

For a particular Lie algebra  $\underline{\alpha}_1 = (1, 0)$ ,  $\underline{\alpha}_2 = (-1, 1)$  and the other positive roots are  $\underline{\alpha}_1 + \underline{\alpha}_2, 2\underline{\alpha}_1 + \underline{\alpha}_2$ . How do these roots correspond to non-zero commutators of  $\{E^+_i\}$ ? Show that for this algebra

$$\begin{aligned} E^-_1|n_1, n_2\rangle &\propto |n_1-2, n_2+1\rangle \quad \text{or} \quad E^-_1|n_1, n_2\rangle = 0, \\ E^-_2|n_1, n_2\rangle &\propto |n_1+2, n_2-2\rangle \quad \text{or} \quad E^-_2|n_1, n_2\rangle = 0. \end{aligned}$$

Find the different states  $|n_1, n_2\rangle$  in the representation space when  $\bar{n}_1 = 2, \bar{n}_2 = 0$ .

[Issues of degeneracy need not be considered.]

**4** For a Lie algebra with a Lie bracket  $[T_a, T_b] = T_c c^c_{ab}$ ,  $a, b, c = 1, \dots, D$  show that the matrices  $(\hat{T}_a)^c_b = c^c_{ab}$  form a representation of the Lie algebra. Let

$$\kappa_{ab} = \text{tr}(\hat{T}_a \hat{T}_b).$$

Explain why  $\kappa_{ab}$  is an invariant tensor satisfying

$$\kappa_{db} c^d_{ca} + \kappa_{ad} c^d_{cb} = 0.$$

Suppose that the Lie algebra has an invariant subalgebra  $\{X_i\}$ , so that  $[T_a, X_i] = X_j c^j_{ai}$  for any  $T_a$ , which is also abelian. Show that then  $\kappa_{ab}$  has an eigenvector with zero eigenvalue.

If  $\kappa_{ab}$  has an inverse  $\kappa^{ab}$  show that for any matrices  $\{t_a\}$  satisfying the Lie algebra

$$[t_a, \kappa^{bc} t_b t_c] = 0.$$

For  $SU(2)$  determine  $\kappa_{ab}$ . What are the expected eigenvalues of  $\kappa^{bc} t_b t_c$  in  $SU(2)$  irreducible representations?