UNIVERSITY OF CAMBRIDGE

MATHEMATICAL TRIPOS Part III

Thursday 8 June, 2006 9 to 11

PAPER 30

INFORMATION AND CODING

*Attempt **THREE** questions.*

*There are **FOUR** questions in total.*

The questions carry equal weight.

STATIONERY REQUIREMENTS
Cover sheet
Treasury Tag
Script paper

SPECIAL REQUIREMENTS
None

You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.

1 (a) Consider two discrete random variables X and Y. Define the conditional entropy $h(X|Y)$, and show that it satisfies

$$h(X|Y) \leqslant h(X),$$

giving necessary and sufficient conditions for equality. You may assume the Gibbs inequality, provided that you state it carefully.

(b) Consider two discrete random variables U and V with corresponding probability mass functions p_U and p_V. For each $\alpha \in [0,1]$, define the mixture random variable $W(\alpha)$ by its mass function

$$p_{W(\alpha)}(x) = \alpha p_U(x) + (1-\alpha) p_V(x).$$

Prove that for all α the entropy of $W(\alpha)$ satisfies:

$$h(W(\alpha)) \geqslant \alpha h(U) + (1-\alpha) h(V).$$

(c) Define $F(\lambda)$ to be the entropy of a Poisson random variable with mean $\lambda > 0$. Show that $F(\lambda)$ is a non-decreasing function of $\lambda > 0$.

2 State the Entropy Power Inequality for n-dimensional random vectors.

Let X be a real-valued random variable with a density and finite differential entropy, and let function $g : \mathbb{R} \to \mathbb{R}$ have strictly positive derivative g' everywhere. Prove that the random variable $g(X)$ has differential entropy satisfying

$$h(g(X)) = h(X) + \mathbb{E} \log_2 g'(X),$$

assuming that $\mathbb{E} \log_2 g'(X)$ is finite.

Let Y_1 and Y_2 be independent, strictly positive random variables with densities. Show that the differential entropy of the product $Y_1 Y_2$ satisfies

$$2^{2h(Y_1 Y_2)} \geqslant \alpha_1 2^{2h(Y_1)} + \alpha_2 2^{2h(Y_2)},$$

where $\log_2(\alpha_1) = 2\mathbb{E} \log_2 Y_2$ and $\log_2(\alpha_2) = 2\mathbb{E} \log_2 Y_1$.

3 (a) Prove the Hamming and Gilbert–Varshamov bounds on the size of a binary code of length N and minimum distance δ, in terms of $v_N(d)$, the volume of an N-dimensional Hamming ball of radius d.

Suppose that the minimum distance is $\lfloor \lambda N \rfloor$ for some fixed $\lambda \in (0, 1/2)$. Describe the asymptotic behaviour of both of the above bounds as $N \to \infty$.

[*You may assume that*

$$\lim_{N\to\infty} \tfrac{1}{N} \log v_N(\lfloor \lambda N \rfloor) = h(\lambda),$$

where h is the binary entropy function].

(b) State Shannon's Second Coding Theorem, giving the capacity of a general memoryless channel. Use this to calculate the capacity of a memoryless binary symmetric channel with error probability p.

(c) Fix $R \in (0, 1)$ and suppose we want to send one of a collection U_N of messages of length N, where the size $|U_N| = 2^{NR}$. The message is transmitted through a memoryless binary symmetric channel with error probability $p < 1/2$, so that we expect about pN errors. According to the asymptotic Gilbert–Varshamov bound of part (a), for which values of p can we correct pN errors, for large N? Why does this give a different answer to the Shannon capacity of part (b)?

4 Prove that the binary code of length 23 generated by the polynomial $g(X) = 1 + X + X^5 + X^6 + X^7 + X^9 + X^{11}$ has minimum distance 7, and is perfect.

[*You may use the BCH theorem without proof provided it is clearly stated, and you may assume that $X^{23} + 1 \equiv (X + 1)g(X)g^{\mathrm{rev}}(X) \bmod 2$, where $g^{\mathrm{rev}}(X) = X^{11}g(1/X)$ is the reversal of $g(X)$.*]

END OF PAPER