

MATHEMATICAL TRIPOS Part III

Friday 30 May 2008 9.00 to 12.00

PAPER 34

ADVANCED PROBABILITY

*Attempt **FOUR** questions.*

*There are **SIX** questions in total.*

The questions carry equal weight.

STATIONERY REQUIREMENTS

*Cover sheet
Treasury Tag
Script paper*

SPECIAL REQUIREMENTS

None

<p>You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.</p>

1 Define the term σ -field. Let $(\mathcal{F}_r : r \in \mathbb{R})$ be a collection of σ -fields of subsets of Ω . Show that $\bigcap_r \mathcal{F}_r$ is a σ -field, but that $\bigcup_r \mathcal{F}_r$ need not be. Show that there exists a smallest σ -field of Ω containing every \mathcal{F}_r .

Let X_1, X_2, \dots be independent random variables on (Ω, \mathcal{F}, P) . Define the tail σ -field of the X_i , and show that every event in the tail σ -field has probability either 0 or 1.

Let $S_n = X_1 + X_2 + \dots + X_n$. Show that the events

$$\left\{ \liminf_{n \rightarrow \infty} S_n / \sqrt{n} \leq -x \right\}, \quad \left\{ \limsup_{n \rightarrow \infty} S_n / \sqrt{n} \geq x \right\}$$

lie in the tail σ -field for every $x \in \mathbb{R}$.

Suppose further that the X_i are symmetric (in that X_i and $-X_i$ have the same distribution), and that there exists $c \in \mathbb{R}$ such that $P(|X_i| \leq c) = 1$ for all i . Show that

$$P\left(|S_n| \leq \frac{1}{2}c \text{ infinitely often}\right) = 1.$$

2 (a) State and prove the two Borel–Cantelli lemmas.

(b) Show that the $N(0, 1)$ distribution function Φ and density function ϕ satisfy

$$1 - \Phi(x) \sim \frac{\phi(x)}{x} \quad \text{as } x \rightarrow \infty.$$

(c) Let X_1, X_2, \dots be independent $N(0, 1)$ random variables on (Ω, \mathcal{F}, P) . Show that

$$P\left(\limsup_{n \rightarrow \infty} \frac{X_n^2}{\log n} = 2\right) = 1.$$

3 Let (Ω, \mathcal{F}, P) be a probability space.

(a) Define the space $L^2 = L^2(\Omega, \mathcal{F}, P)$ and show that it is complete in that, for any Cauchy sequence $X_n \in L^2$ there exists $X \in L^2$ such that $X_n \rightarrow X$ in L^2 .

(b) Define the terms ‘filtration’ and ‘martingale’. State the almost-sure martingale convergence theorem.

(c) Let $(X_n : n \geq 0)$ be a martingale such that $E(X_n^2) \leq M < \infty$ for all n . Show that

$$E\left((X_n - X_m)^2\right) = E(X_n^2) - E(X_m^2)$$

for $m \leq n$, and deduce that X_n converges in L^2 .

4 (a) Let X_1, X_2, \dots be independent integrable random variables with $E(X_i) = 0$ for all i . Show that $S_n = X_1 + X_2 + \dots + X_n$ defines a martingale with respect to the filtration given by

$$\mathcal{F}_n = \sigma(X_1, X_2, \dots, X_n).$$

(b) Suppose further that the X_i are identically distributed and take values in $\{\dots, -2, -1, 0, 1\}$ with $P(X_1 = 1) > 0$. Let $M(t) = E(e^{tX_1})$ and suppose $\tau > 0$. Show that $M(\tau) \in [1, \infty)$, and that

$$Z_n = e^{\tau S_n} / M(\tau)^n$$

is a martingale.

(c) [Continuation] Let $b \in \{1, 2, \dots\}$ and $T = \inf\{n : S_n = b\}$. By considering the non-negative martingale $b - S_{n \wedge T}$ or otherwise, show that $P(T < \infty) = 1$.

(d) [Continuation] Show that the martingale $Z_{n \wedge T}$ is uniformly integrable, and deduce that

$$E(M(\tau)^{-T}) = e^{-\tau b}.$$

(e) Calculate $E(e^{-\alpha T})$ for $\alpha > 0$, in the special case when

$$P(X_1 = 1) = P(X_1 = -1) = \frac{1}{2}.$$

[Any general result which you use should be stated clearly.]

5 (a) Define the term ‘uniformly integrable’. Let Z be integrable and \mathcal{F}_n a filtration. Show that the sequence $X_n = E(Z|\mathcal{F}_n), n \geq 1$, is uniformly integrable.

(b) For $x \in [0, 1)$, define for non-negative integers k, n ,

$$b_n(x) = k2^{-n} \quad \text{if} \quad k2^{-n} \leq x < (k+1)2^{-n}.$$

Let $f : [0, 1] \rightarrow \mathbb{R}$ be integrable, and let U be uniformly distributed on $[0, 1]$. Show that $X_n = E(f(U)|\mathcal{F}_n)$ defines a uniformly integrable martingale with respect to the filtration $\mathcal{F}_n = \sigma(b_n(U))$. Let f_n be the step function on $[0, 1]$ given by

$$f_n(x) = 2^n \int_{b_n(x)}^{b_n(x)+2^{-n}} f(u) du.$$

Show that $f_n(x) \rightarrow f(x)$ for almost every x , and

$$\int_0^1 |f_n(u) - f(u)| du \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty.$$

6 Define a standard Brownian motion $B = (B_t : t \geq 0)$. Give a careful statement of the Strong Markov Property. Set

$$M_t = \sup \{B_s : 0 \leq s \leq t\}.$$

Prove that

$$P(M_t \geq m, B_t \leq x) = P(B_t \geq 2m - x)$$

for $t \geq 0, m > 0$, and $x \leq m$.

Deduce that M_t has the same law as $|B_t|$.

For $x > 0$, let $T_x = \inf\{t : B_t \geq x\}$. Show that T_x has the same law as $(x/B_1)^2$, and calculate the density function of T_x .

END OF PAPER