UNIVERSITY OF CAMBRIDGE

MATHEMATICAL TRIPOS Part III

Monday, 7 June, 2010 9:00 am to 11:00 am

PAPER 31

NONPARAMETRIC STATISTICAL THEORY

*Attempt no more than **THREE** questions.*

*There are **FOUR** questions in total.*

The questions carry equal weight.

STATIONERY REQUIREMENTS
Cover sheet
Treasury Tag
Script paper

SPECIAL REQUIREMENTS
None

You may not start to read the questions printed on the subsequent pages until instructed to do so by the Invigilator.

1

Let $X_1, \ldots, X_n$ be independent and identically distributed random variables with distribution function $F : \mathbb{R} \to [0,1]$. Define the empirical distribution function $F_n : \mathbb{R} \to [0,1]$. Define the (standard) Brownian bridge $\mathbb{G}$ and the F-Brownian bridge process $\mathbb{G}_F$. Carefully state Donsker's central limit theorem for $F_n - F$.

Assuming that F is continuous, use Donsker's central limit theorem to prove that

$$\sqrt{n} \sup_{t \in \mathbb{R}} |F_n(t) - F(t)| \xrightarrow{d} \max_{t \in [0,1]} |\mathbb{G}(t)|$$

$n \to \infty$. [You may use the continuous mapping theorem for convergence in distribution in metric spaces.]

Explain how this result can be used to construct an asymptotic confidence band for F centered at F_n.

2

Let $X_1, \ldots, X_n$ be independent and identically distributed random variables with probability density function $f : \mathbb{R} \to [0, \infty)$. Define the kernel density estimator $f_n^K(x,h)$ of f with bandwidth h.

What is a kernel of order l? Let $\{\phi_m\}_{m \in \mathbb{N}}$ be the orthonormal basis of Legendre polynomials defined by

$$\phi_0(x) := 2^{-1/2}, \qquad \phi_m(x) = \sqrt{\frac{2m+1}{2}} \, \frac{1}{2^m \, m!} \, \frac{d^m}{dx^m} \left[(x^2-1)^m\right]$$

for $x \in [-1,1]$ and $m \in \mathbb{N}$. Using these polynomials (or otherwise), construct a kernel of order l. [You may use standard properties of spaces of polynomials in your answer.]

Suppose f is three times differentiable and that f and $D^3 f$ are bounded functions. Devise a kernel K and a bandwidth h_n depending on n such that for every $x \in \mathbb{R}$

$$E\left|f_n^K(x,h) - f(x)\right| \leqslant C n^{-\frac{3}{7}}$$

for some constant C independent of n. Give an example of a probability density function $f : \mathbb{R} \to [0, \infty)$ for which

$$E\left|f_n^K(x,h) - f(x)\right| \leqslant C n^{-\frac{1}{2}}$$

for every $x \in \mathbb{R}$.

3

If $m : \mathbb{R} \to \mathbb{R}$ is a function for which $\int_{\mathbb{R}} (m(x))^2 \, dx < \infty$, and if ϕ and ψ are the generating functions of a wavelet basis, define carefully the wavelet series and the wavelet projection of m.

Considering random variables Y_i, design points x_i and a regression function m, define the *fixed design regression model*. For design points $x_i = i/n$, define the wavelet regression estimator of m. By analogy to the wavelet thresholding density estimator, argue how one could implement a *thresholded* wavelet regression estimator, and motivate a reasonable choice of thresholds.

4

Let S be a normed space with norm $\| \cdot \|_S$, and let Φ be a real-valued mapping defined on S. Define the notion of Hadamard-differentiability of Φ at a point $s_0 \in S$. Let r_n be real numbers diverging to infinity and let X_n be random variables taking values in S such that $r_n(X_n - s_0)$ converges in distribution to some random variable X taking values in S as $n \to \infty$. Derive the asymptotic distribution of

$$r_n(\Phi(X_n) - \Phi(s_0))$$

as $n \to \infty$. [You may use Skorohod's almost sure representation theorem in the proof, provided it is carefully stated.]

Suppose now $\Phi : L^\infty \to \mathbb{R}$ is Hadamard differentiable on the space L^∞ of bounded real-valued functions defined on $\mathbb{R}$ equipped with the supremum norm $\|f\|_\infty = \sup_{t \in \mathbb{R}} |f(t)|$. Suppose you are given a random sample from some unknown distribution function $F : \mathbb{R} \to [0,1]$. Construct an estimator T_n for $\Phi(F)$ such that $\sqrt{n}(T_n - \Phi(F))$ is stochastically bounded, that is, for every $\varepsilon > 0$ there exists $n_0 := n_0(\varepsilon) \in \mathbb{N}$ and $M := M(\varepsilon)$ finite such that $\Pr(\sqrt{n}(T_n - \Phi(F)) \leqslant M) > 1 - \varepsilon$ for every $n \geqslant n_0$. [You may use Donsker's central limit theorem in the answer.]

END OF PAPER